

ALOCLAIR® PLUS MOUTHWASH - INFORMATION LEAFLET

Aloclair Plus Mouthwash forms a protective film which adheres on the oral cavity affording a mechanical barrier to the affected areas thus relieving the pain caused by exposed nervendings.

Aloclair Plus Mouthwash promotes healing and helps in the management of pain caused by small lesions in the mouth such as those due to aphthous ulcers, aphthous stomatitis and minor lesions; chafing and traumatic ulcers caused by braces and ill-fitting dentures and lesions of surgical origin. It is also indicated for diffuse aphthous ulcers. The hyaluronic acid (high molecular weight) and the Aloe vera in the formulation support the natural healing of damaged tissues.

Aloclair Plus Mouthwash is suitable for the use in adults and children of all ages.

Instructions for use:

Fill the measuring cap provided to the 10ml indicator mark. Rinse around the mouth for at least one minute before expectorating. Use 3-4 times a day, or as needed. For best results, avoid eating or drinking for at least 1 hour after application. Aloclair Plus Mouthwash is safe if ingested.

Ingredients

Aqua, polyvinylpyrrolidone (PVP), maltodextrin, propylene glycol, aloe barbadensis, hydroxy ethylcellulose, potassium sorbate, sodium benzoate, PEG-40 hydrogenated castor oil, disodium EDTA, benzalkonium chloride, sodium hyaluronate, aroma, sodium saccharin, dipotassium glycyrrhizate.

Cautions:

The use of Aloclair Plus Mouthwash is contra-indicated in any patient with a known history of hypersensitivity to any of the ingredients. If you frequently suffer from mouth ulcers seek medica advice

Precautions and warnings:

Do not use if the bottle is damaged or leaking. Store at room temperature, away from direct heat. Keep out of the reach of children

Contents: 60ml Θ and 120ml Θ

Manufacturer:

Alliance Pharma S.r.l. Viale Restelli 5, 20124 Milano, Italy

Aloclair®, Alliance and associated devices are trademarks of the Alliance Pharma PLC group





ET
avity affording exposed nerve
pain caused by comatitis and minor and lesions of acid (high molecular maged tissues.
e mouth for at least st results, avoid h is safe if ingested.
densis, hydroxy- castor oil, n saccharin,
known history of ulcers seek medical
hildren.
arma PLC group
Rev 09/2016 TR1169616 OSIN2FI0134E